

Dr. Patrick McCrea

M.D., M.A.Sc., F.R.C.S.C., General Surgeon

Boardwalk Professional Centre
220 Water Street, Charlottetown, PE, C1A 9M5
Email: info@drpatrickmccrea.com | Web: www.drpatrickmccrea.con
Tel: 902.368.7610 | Fax: 902.368.7611

AFTERNOON COLONOSCOPY AND GASTROSCOPY

This is an instruction sheet for patients scheduled for afternoon colonoscopy *AND* gastroscopy. If you believe you are scheduled otherwise and received this form in error, please call the clinic.

If this is your first visit with Dr. Patrick McCrea, please fill out the Health Questionnaire (attached), and return to the office. The clinic should receive by email (preferred), faxed, or mail at least 1 week in advance. Please watch instructional videos for your procedure on www.drpatrickmccrea.com.

Date:	Registration Time:	Procedure Time:
Location: Endoscopy	y Unit, Queen Elizabeth Hospital, 60 R	iverside Drive, Charlottetown, PE.
days prior to schedu	- •	rea.com to confirm appointment at least 7 vill be cancelled. If you cancel later than this, nay be subject to a cancellation fee.
•	,	a responsible adult take you to and from the arsing to call) and be within 15 minutes of the

REQUIRED BOWEL PREPARATION INSTRUCTIONS:

Supplies Needed: Available without a prescription at any pharmacy.

3 Dulcolax Tablets (Bisacodyl laxative)

Attention:

• Colyte/Klean Prep – 4 litre bottle



Dr. Patrick McCrea

M.D., M.A.Sc., F.R.C.S.C., General Surgeon

Boardwalk Professional Centre
220 Water Street, Charlottetown, PE, C1A 9M5
Email: info@drpatrickmccrea.com | Web: www.drpatrickmccrea.con
Tel: 902.368.7610 | Fax: 902.368.7611

INSTRUCTIONS:

DAY BEFORE APPOINTMENT

8:00am: Start a clear liquid diet. Clear fluids include: Water, clear juices without pulp, chicken broth or consommé soups, light coloured jello without fruits, popsicles, Gatorade, Kool-aid, clear soft drinks, Ovaltine mixed with water. These are just examples. No milk, nothing red or purple. Do not eat seeds or kernels for 2 weeks prior to appointment.

12:00pm (noon): Start Colyte/Klean Prep. Only drink ¾ of this solution. Do not drink too fast. Drink an 8-10 oz glass over 20-30 minutes. It may take several hours to drink all of the prep solution. Leave one litre to drink the following morning.

6:00pm: Take 3 Dulcolax tablets orally.

DAY OF APPOINTMENT

7:00-9:00am: Drink the remaining Colyte. Continue to drink plenty of fluids up to 4 hours before procedure. You will need to have access to a toilet after you take this medication as it can begin to work within 1-3 hours. It is very important to drink a sufficient amount of clear fluids to avoid dehydration. Cramps and diarrhea are expected, so stay close to a toilet. Should you develop nausea, you may take 25mg of Gravol (dimenhydrinate) every 6 hours with water. If you vomit, wait until the nausea subsides and proceed with the prep. A good cleansing is usually achieved even if you vomit, if the prep is continued.

If you are travelling a distance more than 1 hour away, you may select to stay in Charlottetown the night before or contact the GI nurses about an alternate way to take the bowel preparation.

MEDICATIONS/CONDITIONS:

If you are on chemotherapy, please advise Dr. McCrea's office as soon as possible as your colonoscopy will have to be deferred until chemotherapy is completed.

If you are taking blood thinning herbs or medications:

- Ticlid (ticlopidine): Contact our office as soon as possible.
- Pradax (dabigatran): Contact our office as soon as possible.
- Coumadin(Warfarin): Contact our office as soon as possible.
- Plavix (clopidogrel): Contact our office as soon as possible.
- Iron, vitamins, and/or herbal medications: Please discontinue 1 week prior to your appointment.

If you are diabetic, your blood sugar may effected by the bowel prep regime; you should take ½ your normal AM insulin, do not take oral diabetic medications, and check your blood sugar. You should take your blood pressure medication as usual. Please take medications with only a sip of water that morning.

POST COLONOSCOPY AND GASTROSCOPY:



Dr. Patrick McCrea

M.D., M.A.Sc., F.R.C.S.C., General Surgeon

Boardwalk Professional Centre
220 Water Street, Charlottetown, PE, C1A 9M5
Email: info@drpatrickmccrea.com | Web: www.drpatrickmccrea.con
Tel: 902.368.7610 | Fax: 902.368.7611

You must avoid travelling for 7 days post procedure. You are free to eat anything you want afterwards unless you are told otherwise. You should ask Dr. McCrea when you can restart any blood thinning medications; you can resume taking all other medications as usual unless you are told otherwise. Go to the emergency department nearest to you and advise our office if you experience any of the following:

- Chills
- Fever
- Heavy bleeding
- Severe abdominal pain

Please refer to www.drpatrickmccrea.com for videos and more information about your procedure. If there are any questions or concerns regarding these instructions, please contact Lindsay Perry, Administrative Assistant, at Dr. McCrea's Office at info@drpatrickmccrea.com or at 902.368.7610.